

# Implementation of Liu's procedure in MATHEMATICA<sup>®</sup> for use in Relativistic Constitutive Theory

Heiko J. Herrmann

February 2, 2008

## Abstract


The aim of this article is to show, how computer algebra can be used when applying Liu's procedure.

Although MATHEMATICA<sup>®</sup> (a commercial product by Wolfram Research Inc.) is used in this article, it is possible to use other computer algebra systems as well.

## 1 Thermodynamics

### 1.1 Liu's Procedure (A *very* short introduction)

Here a *very* short introduction will be presented, so that the reader will get a glance of what the Liu procedure is. For a detailed discussion of Liu's procedure and balance equations please consult the following literature: [1], [2], [3], [4], [5],[6], [7], [8]



**Proposition 1 (Coleman-Mizel formulation of the 2nd Law)** [9]

If  $\mathcal{Z}$  is no trap, then the following statement is valid for all  $X$ :

$$A(\mathcal{Z}) \cdot X = -B(\mathcal{Z}) \quad (1)$$

$$\Rightarrow \alpha(\mathcal{Z}) \cdot X \geq -\beta(\mathcal{Z}) \quad (2)$$

For all  $X$ , which satisfy the balance equations, the dissipation inequality is fulfilled automatically.

**Proposition 2 (Liu proposition) [1]**

Starting with the Coleman-Mizel formulation of the Second Law one can show that in large state spaces there exist state functions so that the following relations are valid.

$$\begin{aligned}\Lambda(\mathcal{Z}) \cdot A(\mathcal{Z}) &= \alpha(\mathcal{Z}) \\ -\Lambda(\mathcal{Z}) \cdot B(\mathcal{Z}) &\geq -\beta(\mathcal{Z})\end{aligned}$$

This means that there exist restrictions to the constitutive equations.

## 1.2 Example for the Application of Liu's Procedure (for arbitrary non-relativistic balances)

Here a short example for general balances and a state space is given [8]. This example shall demonstrate how the balances on the state space and the matrix formulation is achieved. Therefore two balances and an inequality for arbitrary quantities are considered.

The balances read:

$$\partial_t a_1 + \nabla \cdot (\underline{v} a_1 + b_1) = c_1 \quad (3)$$

$$\partial_t a_2 + \nabla \cdot (\underline{v} a_2 + b_2) = c_2 \quad (4)$$

$$\partial_t \xi + \nabla \cdot (\underline{v} \xi + \gamma) \geq 0 \quad (5)$$

The state space is

$$\mathcal{Z} = \{a_1, a_2, \underline{v}, \partial_t a_1\} \quad (6)$$

This is the domain of for the constitutive functions  $b_1, b_2, \xi, \gamma$ .

Balances on the state space are obtained by using the chain rule when performing the derivatives.

$$\partial_t a_1 + \nabla \cdot (\underline{v} a_1 + b_1(\mathcal{Z})) - c_1 = 0 \quad (7)$$

$$\begin{aligned}& \partial_t a_1 + \nabla \cdot \underline{v} a_1 + \underline{v} \cdot \nabla a_1 + \\ & + \frac{\partial b_1}{\partial a_1} \cdot \nabla a_1 + \frac{\partial b_1}{\partial a_2} \cdot \nabla a_2 + \frac{\partial b_1}{\partial \underline{v}} \cdot \nabla \underline{v} + \frac{\partial b_1}{\partial \partial_t a_1} \cdot \nabla \partial_t a_1 - c_1 = 0\end{aligned} \quad (8)$$

$$\partial_t a_2 + \nabla \cdot (\underline{v} a_2 + b_2(\mathcal{Z})) - c_2 = 0 \quad (9)$$

$$\begin{aligned}& \partial_t a_2 + \nabla \cdot \underline{v} a_2 + \underline{v} \cdot \nabla a_2 + \\ & + \frac{\partial b_2}{\partial a_1} \cdot \nabla a_1 + \frac{\partial b_2}{\partial a_2} \cdot \nabla a_2 + \frac{\partial b_2}{\partial \underline{v}} \cdot \nabla \underline{v} + \frac{\partial b_2}{\partial \partial_t a_1} \cdot \nabla \partial_t a_1 - c_2 = 0\end{aligned} \quad (10)$$

$$\partial_t \xi(\mathcal{Z}) + \nabla \cdot (\underline{v} \xi(\mathcal{Z}) + \gamma(\mathcal{Z})) \geq 0 \quad (11)$$

$$\begin{aligned}& \frac{\partial \xi}{\partial a_1} \cdot \partial_t a_1 + \frac{\partial \xi}{\partial a_2} \cdot \partial_t a_2 + \frac{\partial \xi}{\partial \underline{v}} \cdot \partial_t \underline{v} + \frac{\partial \xi}{\partial \partial_t a_1} \cdot \partial_t \partial_t a_1 + \\ & + \nabla \cdot \underline{v} \xi + \underline{v} \cdot \frac{\partial \xi}{\partial a_1} \nabla a_1 + \underline{v} \cdot \frac{\partial \xi}{\partial a_2} \nabla a_2 + \underline{v} \cdot \frac{\partial \xi}{\partial \underline{v}} \nabla \underline{v} + \underline{v} \cdot \frac{\partial \xi}{\partial \partial_t a_1} \nabla \partial_t a_1 + \\ & + \frac{\partial \gamma}{\partial a_1} \cdot \nabla a_1 + \frac{\partial \gamma}{\partial a_2} \cdot \nabla a_2 + \frac{\partial \gamma}{\partial \underline{v}} \cdot \nabla \underline{v} + \frac{\partial \gamma}{\partial \partial_t a_1} \cdot \nabla \partial_t a_1 \geq 0\end{aligned} \quad (12)$$

As  $\partial_t a_1$  is included in the state space it is not a higher derivative.

Next the balances can be written in matrix formulation.

$$\underline{A}(\mathcal{Z}) \cdot \underline{X} + \underline{B}(\mathcal{Z}) = 0 \quad (13)$$

$$\underline{\alpha}(\mathcal{Z}) \cdot \underline{X} + \beta(\mathcal{Z}) \geq 0 \quad (14)$$

$$((X))^\top = \left( \partial_t a_2 \quad \partial_t \underline{v} \quad \partial_t \partial_t a_1 \quad \nabla a_1 \quad \nabla a_2 \quad \nabla \underline{v} \quad \nabla \partial_t a_1 \right) \quad (15)$$

$$((A)) = \begin{pmatrix} 0 & 0 & 0 & \underline{v} + \frac{\partial b_1}{\partial a_1} & \frac{\partial b_1}{\partial a_2} & a_1 \underline{1} + \frac{\partial b_1}{\partial \underline{v}} & \frac{\partial b_1}{\partial \partial_t a_1} \\ 1 & 0 & 0 & \frac{\partial b_2}{\partial a_1} & \underline{v} + \frac{\partial b_2}{\partial a_2} & a_2 \underline{1} + \frac{\partial b_2}{\partial \underline{v}} & \frac{\partial b_2}{\partial \partial_t a_1} \end{pmatrix} \quad (16)$$

$$((B)) = \begin{pmatrix} \partial_t a_1 - c_1 \\ -c_2 \end{pmatrix} \quad (17)$$

$$((\alpha)) = \left( \frac{\partial \xi}{\partial a_2} \quad \frac{\partial \xi}{\partial \underline{v}} \quad \frac{\partial \xi}{\partial \partial_t a_1} \quad \underline{v} \cdot \frac{\partial \xi}{\partial a_1} + \frac{\partial \gamma}{\partial a_1} \quad \underline{v} \cdot \frac{\partial \xi}{\partial a_2} + \frac{\partial \gamma}{\partial a_2} \quad \xi \underline{1} + \underline{v} \cdot \frac{\partial \xi}{\partial \underline{v}} + \frac{\partial \gamma}{\partial \underline{v}} \quad \underline{v} \cdot \frac{\partial \xi}{\partial \partial_t a_1} + \frac{\partial \gamma}{\partial \partial_t a_1} \right) \quad (18)$$

$$((\beta)) = \left( \frac{\partial \xi}{\partial a_1} \cdot \partial_t a_1 \right) \quad (19)$$

Next the Liu equations can be written down

$$\underline{\Lambda}(\mathcal{Z}) \cdot \underline{A}(\mathcal{Z}) = \underline{\alpha}(\mathcal{Z}) \quad (20)$$

$$-\underline{\Lambda}(\mathcal{Z}) \cdot \underline{B}(\mathcal{Z}) \geq -\beta(\mathcal{Z}) \quad (21)$$

$$\Lambda_2 = \frac{\partial \xi}{\partial a_2} \quad (22)$$

$$0 = \frac{\partial \xi}{\partial \underline{v}} \quad (23)$$

$$0 = \frac{\partial \xi}{\partial \partial_t a_1} \quad (24)$$

$$\Lambda_1 \left( \underline{v} + \frac{\partial b_1}{\partial a_1} \right) + \Lambda_2 \left( \frac{\partial b_2}{\partial a_1} \right) = \underline{v} \cdot \frac{\partial \xi}{\partial a_1} + \frac{\partial \gamma}{\partial a_1} \quad (25)$$

$$\Lambda_1 \frac{\partial b_1}{\partial a_2} + \Lambda_2 \left( \underline{v} + \frac{\partial b_2}{\partial a_2} \right) = \underline{v} \cdot \frac{\partial \xi}{\partial a_2} + \frac{\partial \gamma}{\partial a_2} \quad (26)$$

$$\Lambda_1 \left( a_1 \underline{1} + \frac{\partial b_1}{\partial \underline{v}} \right) + \Lambda_2 \left( a_2 \underline{1} + \frac{\partial b_2}{\partial \underline{v}} \right) = \xi + \underline{v} \cdot \frac{\partial \xi}{\partial \underline{v}} + \frac{\partial \gamma}{\partial \underline{v}} \quad (27)$$

$$\Lambda_1 \frac{\partial b_1}{\partial \partial_t a_1} + \Lambda_2 \frac{\partial b_2}{\partial \partial_t a_1} = \underline{v} \cdot \frac{\partial \xi}{\partial \partial_t a_1} + \frac{\partial \gamma}{\partial \partial_t a_1} \quad (28)$$

The residual inequality reads

$$\Lambda_1 (\partial_t a_1 - c_1) - \Lambda_2 c_2 \leq \frac{\partial \xi}{\partial a_1} \cdot \partial_t a_1 \quad (29)$$

Because of the principle of *material frame indifference* all derivatives with respect to  $\underline{v}$  shall be set equal to zero:  $\frac{\partial}{\partial \underline{v}} \stackrel{!}{=} 0$ . The *material frame indifference* means that material laws shall be independent of the velocity of the observer. Using this the Liu equations read:

$$\Lambda_2 = \frac{\partial \xi}{\partial a_2} \quad (30)$$

$$0 = \frac{\partial \xi}{\partial \underline{v}} \quad (31)$$

$$0 = \frac{\partial \xi}{\partial \partial_t a_1} \quad (32)$$

$$\Lambda_1 \left( \underline{v} + \frac{\partial b_1}{\partial a_1} \right) + \Lambda_2 \left( \frac{\partial b_2}{\partial a_1} \right) = \underline{v} \cdot \frac{\partial \xi}{\partial a_1} + \frac{\partial \gamma}{\partial a_1} \quad (33)$$

$$\Lambda_1 \frac{\partial b_1}{\partial a_2} + \Lambda_2 \left( \underline{v} + \frac{\partial b_2}{\partial a_2} \right) = \underline{v} \cdot \frac{\partial \xi}{\partial a_2} + \frac{\partial \gamma}{\partial a_2} \quad (34)$$

$$\Lambda_1 (a_1 \underline{1}) + \Lambda_2 (a_2 \underline{1}) = \xi \quad (35)$$

$$\Lambda_1 \frac{\partial b_1}{\partial \partial_t a_1} + \Lambda_2 \frac{\partial b_2}{\partial \partial_t a_1} = \underline{v} \cdot \frac{\partial \xi}{\partial \partial_t a_1} + \frac{\partial \gamma}{\partial \partial_t a_1} \quad (36)$$

The residual inequality remains unchanged

$$\Lambda_1 (\partial_t a_1 - c_1) - \Lambda_2 c_2 \leq \frac{\partial \xi}{\partial a_1} \cdot \partial_t a_1 \quad (37)$$

From these equations the Lagrange parameters  $\underline{\Lambda}(\underline{\mathcal{Z}})$  can be determined.

$$\Lambda_2 = \frac{\partial \xi}{\partial a_2} \quad (38)$$

$$\Lambda_1 = \left( \xi - \frac{\partial \xi}{\partial a_2} (a_2 \underline{1}) \right) (a_1 \underline{1})^{-1} \quad (39)$$

After inserting the Lagrange parameters into Liu's equations the restrictions to constitutive equations are given

$$0 = \frac{\partial \xi}{\partial \underline{v}} \quad (40)$$

$$0 = \frac{\partial \xi}{\partial \partial_t a_1} \quad (41)$$

$$\left( \xi - \frac{\partial \xi}{\partial a_2} (a_2 \underline{1}) \right) (a_1 \underline{1})^{-1} \left( \underline{v} + \frac{\partial b_1}{\partial a_1} \right) + \frac{\partial \xi}{\partial a_2} \left( \frac{\partial b_2}{\partial a_1} \right) = \underline{v} \cdot \frac{\partial \xi}{\partial a_1} + \frac{\partial \gamma}{\partial a_1} \quad (42)$$

$$\left( \xi - \frac{\partial \xi}{\partial a_2} (a_2 \underline{1}) \right) (a_1 \underline{1})^{-1} \frac{\partial b_1}{\partial a_2} + \frac{\partial \xi}{\partial a_2} \left( \underline{v} + \frac{\partial b_2}{\partial a_2} \right) = \underline{v} \cdot \frac{\partial \xi}{\partial a_2} + \frac{\partial \gamma}{\partial a_2} \quad (43)$$

$$\left( \xi - \frac{\partial \xi}{\partial a_2} (a_2 \underline{1}) \right) (a_1 \underline{1})^{-1} \frac{\partial b_1}{\partial \partial_t a_1} + \frac{\partial \xi}{\partial a_2} \frac{\partial b_2}{\partial \partial_t a_1} = \underbrace{\underline{v} \cdot \frac{\partial \xi}{\partial \partial_t a_1}}_{=0} + \frac{\partial \gamma}{\partial \partial_t a_1} \quad (44)$$

The entropy production is given by the residual inequality

$$\sigma := - \left( \underline{v} \cdot \frac{\partial \xi}{\partial a_2} \right) \left( \frac{\partial b_1}{\partial a_2} \right)^{-1} (\partial_t a_1 - c_1) + \frac{\partial \xi}{\partial a_1} \geq 0 \quad (45)$$

This example was meant to demonstrate how the Liu equations and the residual inequality are achieved. This example shows also that the entropy production is independent of the highest derivatives included in the state space (this is a general property of the entropy density; see also [10]).

## 2 Application of Liu's Procedure using Mathematica

For the correct handling of indices I will use the package "Ricci.m" by John M. Lee, which can be found on the MathSource server.

```
<<Ricci'
```

### 2.1 Declaration of Tensors and State Space

First it is necessary to define a bundle and declare the letters used for indices.

```
DefineBundle[fiber2,4,g,{i,j,k,l,m,n,o,r,y}]
```

Next the tensors have to be defined. First I will declare the state space variables ( $z_0, \dots, z_5$ ).

$$\mathcal{Z} = \{n_0, u, p, e, spa, spv\}$$

```
(* z0=n0 *) DefineTensor[n0,0] (* particle number density *)
(* z1=u *) DefineTensor[u,1] (* four velocity *)
(* z2=p *) DefineTensor[p,1] (* momentum flux *)
(* z3=e *) DefineTensor[e,0] (* energy *)
(* z4=spa *) DefineTensor[spa,2, Symmetries -> Skew] (* spin density *)
(* z5=spv *) DefineTensor[spv,1] (* vector spin density *)
```

Then the constitutive functions are declared:

```
DefineTensor[t,2] (* stress tensor *)
DefineTensor[q,1] (* heat flux *)
DefineTensor[scs,3] (* couple stress *)
DefineTensor[spb,2] (* 2-couple stress *)
DefineTensor[s,0] (* entropy *)
DefineTensor[se,1] (* entropy flux *)
DefineTensor[f,1] (* external force *)
```

## 2.2 Balance Equations

The next step is to write down the balance equations, which are for technical reasons resolved to zero, and the right hand side ( $= 0$ ) is not written down:

Balance of particle number density:

$$\partial_l (nu^l) = 0$$

$$D[n0[x] u[U[1]] [x], x]$$

Balance of energy momentum

$$\partial_l (t^{il} + \frac{1}{c^2} p^i u^l + \frac{1}{c^2} u^i q^l + \frac{1}{c^4} e u^i u^l) - f^i = 0$$

$$D[t[U[i], U[1]] [x] + 1/c^2 p[U[i]] [x] u[U[1]] [x] + 1/c^2 u[U[i]] [x] q[U[1]] [x] + 1/c^4 e[x] u[U[i]] [x] u[U[1]] [x], x] - f[U[i]]$$

Balance of spin

$$\partial_l \left( s c s^{ijl} + \frac{1}{c^2} s p a^{ij} u^l + \frac{1}{c^2} u^{[i} s p b^{j]l} + \frac{1}{c^4} u^{[i} s p v^{j]} u^l \right) - \left( t^{[ij]} + \frac{1}{c^2} p^{[i} u^{j]} + \frac{1}{c^2} u^{[i} q^{j]} \right) = 0$$

$$D[s c s[U[i], U[j], U[1]] [x] + 1/c^2 s p a[U[i], U[j]] u[U[1]] + 1/(2 c^2) u[U[i]] s p b[U[j], U[1]] [x] - 1/(2 c^2) u[U[j]] s p b[U[i], U[1]] [x] + 1/(2 c^4) u[U[i]] s p v[U[j]] u[U[1]] - 1/(2 c^4) u[U[j]] s p v[U[i]] u[U[1]], x] - 1/2 (t[U[i], U[j]] + 1/c^2 p[U[i]] u[U[j]] + 1/c^2 u[U[i]] q[U[j]] - t[U[j], U[i]] + 1/c^2 p[U[j]] u[U[i]] + 1/c^2 u[U[j]] q[U[i]])$$

Balance of entropy

$$\partial_l \left( s u^l + \frac{1}{c^2} s e^l \right) \geq 0$$

$$D[s [x] u[U[1]] [x] + 1/c^2 s e[U[1]] [x], x]$$

## 2.3 Balances on State Space

As the constitutive functions are defined on the state space, the derivatives have to be performed by use of the chain rule. The equations are "labeled" for further use.

$$\partial_l (nu^l) = 0$$

$$TB = D[n_0[x] \ u[U[1]] [x], x]$$

$$\partial_l \left( t^{il}(\mathcal{Z}) + \frac{1}{c^2} p^i u^l + \frac{1}{c^2} u^i q^l(\mathcal{Z}) + \frac{1}{c^4} e u^i u^l \right) - f^i = 0$$

$$\text{ImpB} = \backslash [\text{Kappa}]$$

$$\begin{aligned} & D[t[U[i], U[1]] [z_0[x], z_1[x], z_2[x], z_3[x], z_4[x], z_5[x]] + \\ & \quad 1/c^2 \ p[U[i]] [x] u[U[1]] [x] + \\ & \quad 1/c^2 \ u[U[i]] [x] q[U[1]] [z_0[x], z_1[x], z_2[x], z_3[x], z_4[x], z_5[x]] + \\ & \quad 1/c^4 \ e[x] u[U[i]] [x] u[U[1]] [x], x] - f[U[i]] \end{aligned}$$

$$\partial_l \left( s c s^{ijl}(\mathcal{Z}) + \frac{1}{c^2} u^{[i} s p b^{j]l}(\mathcal{Z}) + \frac{1}{c^4} u^{[i} s p v^{j]} u^l \right) - \left( t^{[ij]} + \frac{1}{c^2} p^{[i} u^{j]} + \frac{1}{c^2} u^{[i} q^{j]} \right) = 0$$

$$\begin{aligned} \text{SpinB} = & D[s c s[U[i], U[j], U[1]] [z_0[x], z_1[x], z_2[x], z_3[x], z_4[x], z_5[x]] + \\ & \quad 1/c^2 \ s p a[U[i], U[j]] \ u[U[1]] + \\ & \quad 1/(2 \ c^2) u[U[i]] \ s p b[U[j], U[1]] [z_0[x], z_1[x], z_2[x], z_3[x], z_4[x], z_5[x]] - \\ & \quad 1/(2 \ c^2) u[U[j]] \ s p b[U[i], U[1]] [z_0[x], z_1[x], z_2[x], z_3[x], z_4[x], z_5[x]] + \\ & \quad 1/(2 \ c^4) u[U[i]] \ s p v[U[j]] \ u[U[1]] - \\ & \quad 1/(2 \ c^4) u[U[j]] \ s p v[U[i]] \ u[U[1]], x] - \\ & \quad 1/2 \ (t[U[i], U[j]] + 1/c^2 \ p[U[i]] \ u[U[j]] + 1/c^2 \ u[U[i]] q[U[j]] - \\ & \quad \quad t[U[j], U[i]] + 1/c^2 \ p[U[j]] \ u[U[i]] + 1/c^2 \ u[U[j]] q[U[i]]) \end{aligned}$$

$$\partial_l \left( s(\mathcal{Z}) u^l + \frac{1}{c^2} s e^l(\mathcal{Z}) \right) \geq 0$$

$$\text{EntropieB} =$$

$$\begin{aligned} & D[s \ [z_0[x], z_1[x], z_2[x], z_3[x], z_4[x], z_5[x]] u[U[1]] [x] + \\ & \quad 1/c^2 \ s e[U[1]] [z_0[x], z_1[x], z_2[x], z_3[x], z_4[x], z_5[x]], x] \end{aligned}$$

## 2.4 Matrix Formulation

Now the components of the matrices for the matrix formulation of the balances have to be collected.

Matrix of higher derivatives

```
X0:=z0'[x]
X1:=z1'[x]
X2:=z2'[x]
X3:=z3'[x]
X4:=z4'[x]
X5:=z5'[x]
```

Then the higher derivatives have to be set correctly by hand.

```
(z0)'[x_] := n0[L[1]]
(z1)'[x_] := u[U[n]] [L[1]]
(z2)'[x_] := p[U[n]] [L[1]]
(z3)'[x_] := e[L[1]]
(z4)'[x_] := spa[U[n], U[m]] [L[1]]
(z5)'[x_] := spv[U[n]] [L[1]]
(n0)'[x_] := n0[L[1]]
(u[U[i]])'[x_] := u[U[i]] [L[1]]
(u[U[j]])'[x_] := u[U[j]] [L[1]]
(u[U[1]])'[x_] := u[U[1]] [L[1]]
(p[U[i]])'[x_] := p[U[i]] [L[1]]
(p[U[j]])'[x_] := p[U[j]] [L[1]]
(e)'[x_] := e[L[1]]
```

Then the coefficient matrix

```
A00=Coefficient[TB,z0'[x]]
A01=Coefficient[TB,z1'[x]]+g[L[n],U[1]] Coefficient[TB,(u[U[1]])'[x]]
A02=Coefficient[TB,z2'[x]]
A03=Coefficient[TB,z3'[x]]
A04=Coefficient[TB,z4'[x]]
A05=Coefficient[TB,z5'[x]]
A10=Coefficient[ImpB,z0'[x]]
A11=Coefficient[ImpB,z1'[x]]+g[L[n],U[i]] Coefficient[ImpB,(u[U[i]])'[x]]+
    g[L[n],U[1]] Coefficient[ImpB,(u[U[1]])'[x]]
A12=Coefficient[ImpB,z2'[x]]+g[L[n],U[i]] Coefficient[ImpB,(p[U[i]])'[x]]
A13=Coefficient[ImpB,z3'[x]]
A14=Coefficient[ImpB,z4'[x]]
A15=Coefficient[ImpB,z5'[x]]
A20=Coefficient[SpinB,z0'[x]]
A21=Coefficient[SpinB,z1'[x]]+g[L[n],U[i]] Coefficient[SpinB,(u[U[i]])'[x]] +
    g[L[n],U[j]] Coefficient[SpinB,(u[U[j]])'[x]]+
    g[L[n],U[1]] Coefficient[SpinB,(u[U[1]])'[x]]
A22=Coefficient[SpinB,z2'[x]]+g[L[n],U[i]] Coefficient[SpinB,(p[U[i]])'[x]]+
```



```

g[L[n],U[j]] Coefficient[SpinB,(p[U[j]])'[x]]
A23=Coefficient[SpinB,z3'[x]]
A24=Coefficient[SpinB,z4'[x]]
A25=Coefficient[SpinB,z5'[x]]

```

Coefficient matrix of dissipation inequality

```

\[Alpha]0=Coefficient[EntropieB,z0'[x]]
\[Alpha]1=
  Coefficient[EntropieB,z1'[x]]+
  g[L[n],U[1]] Coefficient[EntropieB,(u[U[1]])'[x]]
\[Alpha]2=Coefficient[EntropieB,z2'[x]]
\[Alpha]3=Coefficient[EntropieB,z3'[x]]
\[Alpha]4=Coefficient[EntropieB,z4'[x]]
\[Alpha]5=Coefficient[EntropieB,z5'[x]]

```

The residual matrix

```

B0=Simplify[
  TB - (A00 z0'[x] + A01 z1'[x]+A02 z2'[x]+A03 z3'[x]+A04 z4'[x]+
    A05 z5'[x])]
B1=Simplify[
  ImpB - (A10 z0'[x] + A11 z1'[x]+A12 z2'[x]+A13 z3'[x]+A14 z4'[x]+
    A15 z5'[x])]
B2= Simplify[
  SpinB - (A20 z0'[x] + A21 z1'[x]+A22 z2'[x]+A23 z3'[x]+A24 z4'[x]+
    A25 z5'[x])]

\[Beta]= Simplify[
  EntropieB - (
    \[Alpha]0 z0'[x] + \[Alpha]1 z1'[x]+\[Alpha]2 z2'[x]+\[Alpha]3 z3'[x]+
    \[Alpha]4 z4'[x]+\[Alpha]5 z5'[x])]

```

## 2.5 Liu Relations

Now Liu's proposition (based on Farkash's lemma) can be applied. The result are the Liu equations

$$\begin{aligned} \text{liu1} &= \lambda_0 A_{00} + \lambda_1 A_{10} + \lambda_2 A_{20} - \alpha_0 \\ \text{liu2} &= \lambda_0 A_{01} + \lambda_1 A_{11} + \lambda_2 A_{21} - \alpha_1 \\ \text{liu3} &= \lambda_0 A_{02} + \lambda_1 A_{12} + \lambda_2 A_{22} - \alpha_2 \\ \text{liu4} &= \lambda_0 A_{03} + \lambda_1 A_{13} + \lambda_2 A_{23} - \alpha_3 \\ \text{liu5} &= \lambda_0 A_{04} + \lambda_1 A_{14} + \lambda_2 A_{24} - \alpha_4 \\ \text{liu6} &= \lambda_0 A_{05} + \lambda_1 A_{15} + \lambda_2 A_{25} - \alpha_5 \end{aligned}$$

and the residual inequality

$$\lambda_0 B_0 + \lambda_1 B_1 + \lambda_2 B_2 - \beta$$

The Lagrange parameters can be determined

$$\begin{aligned} \text{lsg} &= \text{Solve}[\{\text{liu1}==0, \text{liu2}==0, \text{liu5}==0\}, \{\lambda_0, \lambda_1, \lambda_2\}]; \\ \text{l0b} &= \lambda_0 /. \text{lsg}[[1]]; \\ \text{l1b} &= \lambda_1 /. \text{lsg}[[1]]; \\ \text{l2b} &= \lambda_2 /. \text{lsg}[[1]]; \\ \text{l1b} &= \lambda_1 /. \text{lsg}[[1]]; \\ \lambda_0 &= \text{l0b} \\ \lambda_1 &= \text{l1b} \\ \lambda_2 &= \text{l2b} \end{aligned}$$

Now one can insert the Lagrange parameters into the Liu equations and one gets the restrictions to the constitutive equations. The residual inequality determines the entropy production density.

# References

- [1] I-Shih Liu. Method of Lagrange Multipliers for Exploitation of the Entropy Principle. *Arch. Rat. Mech. Anal.*, 46:131, 1972.
- [2] W. Muschik, C. Papenfuß, and H. Ehrentraut. *Concepts of Continuum Thermodynamics*. Technische Universität Berlin und Kielce University of Technology, 1996.
- [3] W. Muschik. *Aspects of Non-Equilibrium Thermodynamics*. World Scientific, Singapore, 1990.
- [4] W. Muschik, C. Papenfuß, and H. Ehrentraut. A sketch of continuum thermodynamics. *J. Non-Newtonian Fluid Mech.*, 96:255–290, 2001.
- [5] H. Herrmann, W. Muschik, C. Papenfuß, and G. Rückner. LIU-technique for exploiting the dissipation-inequality in case of constrained state space variables. In *Trends in Application of Mathematics to Mechanics*. Chapman & Hall/CRC, 2000.
- [6] H. Herrmann, W. Muschik, C. Papenfuß, and G. Rückner. Exploiting the dissipation-inequality in case of constrained state space variables. In *Proceedings of the International Symposium on Trends in Continuum Physics TRECOP'98*, Singapore, August 17 - 20 1998. World Scientific.
- [7] W. Muschik and H. Ehrentraut. An amendment to the second law. *J. Non-Equilib. Thermodyn.*, 21:175, 1996.
- [8] Heiko Herrmann. *Non-equilibrium Thermodynamics in Geometrized Theories of Gravitation with Curvature and Torsion*. dissertaion.de Verlag im Internet, Dissertation, Fakultät II, TU Berlin, D83, 2003. ISBN 3-89825-662-6.
- [9] B.D. Coleman and V.J. Mizel. Existence of caloric equations of state in thermodynamics. *J. Chem. Phys.*, 40:1116–1125, 1964.
- [10] W. Muschik and R. Dominguez-Cascante. On extended thermodynamics of discrete systems. *Physica A*, 233:523–550, 1996.